

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Then a-A=water in the cask.

A+b=quantity of wine in cask before the (m+1)th draught since b gallons of wine are added.

$$A+b-[(A/n)+(b/n)]+b=A\left(\frac{n-1}{n}\right)+b\left(\frac{n-1}{n}\right)$$
 =quantity of wine be-

fore the (m+2)th draught.

$$A\left(\frac{n-1}{n}\right) + b\left(\frac{2n-1}{n}\right) - A\left(\frac{n-1}{n^2}\right) - b\left(\frac{2n-1}{n^2}\right) + b = A\left(\frac{n-1}{n}\right)^2 - b\left(\frac{3n^2 - 3n + 1}{n^2}\right)$$

=quantity of wine before the (m+3)th draught.

$$\therefore A \left( \frac{n-1}{n} \right)^{p} + b \left( p n^{p-1} - \frac{p(p-1)}{1 \cdot 2} n^{p-2} + \dots \right) + b$$

$$= A \left( \frac{n-1}{n} \right)^{p} + b \left( \frac{n^{p} - (n-1)^{p}}{n^{p}} \right)$$

=quantity of wine left after 
$$(m+p)$$
th draught= $a\left(\frac{n-1}{n}\right)^{m+p}+b\left(\frac{n^p-(n-1)^p}{n^p}\right)$ 

In the present case, a=10, b=1,  $1/m=\frac{1}{10}$ , m=5, and p=5. Hence, sub-

stituting, we have 
$$10 \left[ \frac{10-1}{10} \right]^{10} + 1 \cdot \left[ \frac{10^5 - (10-1)^5}{10^5} \right] = 7.581884401$$
 gallons,

the quantity of wine left after putting in the last gallon of wine, and, therefore, 2.418115599 gallons—quantity of water in the cask.

## GEOMETRY.

Conducted by B. F. FINKEL. Springfield, Mo. All contributions to this department should be sent to him.

## SOLUTIONS OF PROBLEMS.

71. Proposed by ROBERT J. ALEY, A. M., Ph. D., Professor of Mathematics, Indiana University, Bloomington, Indiana.

Prove by pure geometry: A perpendicular at the middle point,  $M_a$ , of the side BC of the triangle ABC meets the circumcircle in A'. On this perpendicular A'' and A''' are taken so that  $M_aA''=M_aA'$  and A''A'''=AH. (H is the orthocenter of triangle ABC.) Prove that A''' is on the circumcircle.


Solution by F. M. McGAW, A. M., Professor of Mathematics, Bordentown Military Institute, Bordentown, New Jersey, and the PROPOSER.

Let  $M_aA_1=M_aA_2$ ,  $A_2A_3=AH$ , to prove  $A_3$  on the circumference of the circle. Since  $A_2A_3$  is a line through M, the center of the circle, the proposition is in effect to prove  $A_3$  one extremity of the diameter through  $M_a$ .

By the conditions  $AH=A_2A_3$ , and is parallel to it, therefore  $AHA_3A_2$  is a parallelogram.

Also triangles BHA and  $M_aMM_b$  are similar, hence since  $2M_aM_b=AB$ , we have  $AH=2MM_a$ .

Therefore, 
$$A_1A_3 = A_2A_3 + A_2M_a + M_aA_1$$
  
 $=AH + 2M_aA_1$   
 $=2M_aM + 2M_aA_1$   
 $=2(MA_1)=2r$ , hence  $A_3$  is extremity of diameter.



Q. E. D.

Also solved by CHAS. C. CROSS, and J. W. SCROGGS. Mr. Cross furnished two different solutions.

72. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics, Columbian University, Washington, D. C.

If a line with its extremities upon two curves move in any manner whatever, (the line may vary in length), and P a point upon the line which divides it in the ratio m:n describe a curve, the area of this curve will be given by the formula—

$$A = \frac{(m^2 + nm)A_1 + (n^2 + mn)A_2 - mnA_3}{(m+n)^2}.$$

No solution of this problem has been received.

73. Proposed by ROBERT J. ALEY, A. M., Ph. D., Professor of Mathematics, Indiana University, Bloomington, Indiana.

Prove by pure geometry: (1) A', B', and C' are the middle points of the arcs BC, CA, and AB respectively. With these points as centers, circles are described passing through B and C, C and A, and A and B respectively. Prove that these circles intersect in O, the center of the incircle of the triangle ABC; (2) that O, the center of the incircle, is Nagel's point of the triangle formed by joining the middle points of the sides.

Solution by CHARLES C. CROSS, Laytonsville, Maryland, and the PROPOSER.

(1) AO cuts the circumcircle at A', for AO bisects angle A and also its subtending arc.  $\angle OBA' = \frac{1}{2}(A+B)$ .

 $\angle BOA' = \frac{1}{2}(A+B)$  for it is exterior angle to triangle BOA.

 $\therefore$  triangle A'BO is isosceles.

A'B=A'O. By similar reasoning it is proved that B'A=B'O and C'A=C'O.

- ... The circles intersect in O.
- (2) It is a well known property of Nagel's point that AQ and  $OM_a$ , BQ and  $OM_b$ , CQ and  $OM_c$  are respectively parallel.

